

# **Markscheme**

## **Specimen paper**

### **Mathematics: applications and interpretation**

#### **Higher level**

#### **Paper 2**

## Instructions to Examiners

### Abbreviations

- M** Marks awarded for attempting to use a correct **Method**.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- R** Marks awarded for clear **Reasoning**.
- AG** Answer given in the question and so no marks are awarded.

### Using the markscheme

#### 1 General

*Award marks using the annotations as noted in the markscheme eg **M1**, **A2**.*

#### 2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, e.g. **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **A1** for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies **M2**, **N3**, etc., do **not** split the marks, unless there is a note.
- Once a correct answer to a question or part-question is seen, ignore further correct working. However, if further working indicates a lack of mathematical understanding do not award the final **A1**. An exception to this may be in numerical answers, where a correct exact value is followed by an incorrect decimal. However, if the incorrect decimal is carried through to a subsequent part, and correct **FT** working shown, award **FT** marks as appropriate but do not award the final **A1** in that part.

#### Examples

	Correct answer seen	Further working seen	Action
1.	$8\sqrt{2}$	5.65685... (incorrect decimal value)	Award the final <b>A1</b> (ignore the further working)
2.	$\frac{1}{4} \sin 4x$	$\sin x$	Do not award the final <b>A1</b>
3.	$\log a - \log b$	$\log(a - b)$	Do not award the final <b>A1</b>

### 3 Implied marks

*Implied marks appear in **brackets e.g. (M1)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.*

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

### 4 Follow through marks (only applied after an error is made)

*Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) or subpart(s). Usually, to award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if the only marks awarded in a subpart are for the answer (i.e. there is no working expected), then **FT** marks should be awarded if appropriate.*

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (e.g. probability greater than 1, use of  $r > 1$  for the sum of an infinite GP,  $\sin \theta = 1.5$ , non integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word “their” in a description, to indicate that candidates may be using an incorrect value.
- Exceptions to this rule will be explicitly noted on the markscheme.
- If a candidate makes an error in one part, but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the question says hence. It is often possible to use a different approach in subsequent parts that does not depend on the answer to previous parts.

### 5 Mis-read

*If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). Apply a **MR** penalty of 1 mark to that question*

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. probability greater than 1,  $\sin \theta = 1.5$ , non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Mis-copying of candidates’ own work does **not** constitute a misread, it is an error.
- The **MR** penalty can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

## 6 Alternative methods

*Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme*

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.

## 7 Alternative forms

*Unless the question specifies otherwise, **accept** equivalent forms.*

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

## 8 Accuracy of Answers

*If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. There are two types of accuracy errors, and the final answer mark should not be awarded if these errors occur.*

- **Rounding errors:** only applies to final answers not to intermediate steps.
- **Level of accuracy:** when this is not specified in the question the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

## 9 Calculators

*A GDC is required for this examination, but calculators with symbolic manipulation features/ CAS functionality are not allowed.*

### Calculator notation

The subject guide says:

*Students must always use correct mathematical notation, not calculator notation.*

Do **not** accept final answers written using calculator notation. However, do not penalize the use of calculator notation in the working.

1. (a)  $2(8 \times 4 + 3 \times 4 + 3 \times 8)$  **M1**  
 $= 136 \text{ (cm}^2\text{)}$  **A1**  
**[2 marks]**

(b)  $\sqrt{8^2 + 4^2 + 3^2}$  **M1**  
 $(AG \Rightarrow) 9.43 \text{ (cm)} (9.4339\dots, \sqrt{89})$  **A1**  
**[2 marks]**

(c)  $-2x + 220 = 0$  **M1**  
 $x = 110$  **A1**  
 $110\,000 \text{ (boxes)}$  **A1**  
**[3 marks]**

(d)  $P(x) = \int -2x + 220 \, dx$  **M1**

**Note:** Award **M1** for evidence of integration.

$P(x) = -x^2 + 220x + c$  **A1A1**

**Note:** Award **A1** for either  $-x^2$  or  $220x$  award **A1** for both correct terms and constant of integration.

$1700 = -(20)^2 + 220(20) + c$  **M1**

$c = -2300$

$P(x) = -x^2 + 220x - 2300$  **A1**  
**[5 marks]**

(e)  $-x^2 + 220x - 2300 = 0$  **M1**  
 $x = 11.005$  **A1**  
 $11\,006 \text{ (boxes)}$  **A1**

**Note:** Award **M1** for their  $P(x) = 0$ , award **A1** for their correct solution to  $x$ . Award the final **A1** for expressing their solution to the minimum number of boxes. Do not accept 11 005, the nearest integer, nor 11 000, the answer expressed to 3 significant figures, as these will not satisfy the demand of the question.

**[3 marks]**

**Total [15 marks]**

2. (a) (i)  $P(Y) = 0.8 \times 0.1 + 0.2 \times 0.3$  **M1**  
 $= 0.14$  **A1**

(ii)  $P(\text{Star} | Y) = \frac{0.8 \times 0.1}{0.14}$  **M1**  
 $= 0.571 \left( \frac{4}{7}, 0.571428... \right)$  **A1**

**[4 marks]**

(b) the colours of the sweets are distributed according to manufacturer specifications **A1**  
**[1 mark]**

(c)

Colour	Brown	Red	Green	Orange	Yellow	Purple
Expected Frequency	12	20	16	16	8	8

**A2**

**Note:** Award **A2** for all 6 correct expected values,  
**A1** for 4 or 5 correct values, **A0** otherwise.

**[2 marks]**

(d) 5 **A1**  
**[1 mark]**

(e) 0.469 (0.4688117...) **A2**  
**[2 marks]**

(f) since  $0.469 > 0.05$  **R1**  
fail to reject the null hypothesis. There is insufficient evidence to  
reject the manufacturer's specifications **A1**

**Note:** Award **R1** for a correct comparison of their correct  $p$ -value to the  
test level, award **A1** for the correct result from that comparison.  
Do not award **R0A1**.

**[2 marks]**

**Total [12 marks]**

3. (a) (i)  $N = 24$   
 $I\% = 14$   
 $PV = -14000$   
 $FV = 0$   
 $P/Y = 4$   
 $C/Y = 4$  **(M1)(A1)**

**Note:** Award **M1** for an attempt to use a financial app in their technology, award **A1** for all entries correct. Accept  $PV = 14000$ .

(€)871.82 **A1**

- (ii)  $4 \times 6 \times 871.82$  **(M1)**  
 (€)20923.68 **A1**

- (iii)  $20923.68 - 14000$  **(M1)**  
 (€)6923.68 **A1**

**[7 marks]**

- (b) (i)  $0.9 \times 14000 (= 14000 - 0.10 \times 14000)$  **M1**  
 (€)12600.00 **A1**

- (ii)  $N = 72$   
 $PV = 12600$   
 $PMT = -250$   
 $FV = 0$   
 $P/Y = 12$   
 $C/Y = 12$  **(M1)(A1)**

**Note:** Award **M1** for an attempt to use a financial app in their technology, award **A1** for all entries correct. Accept  $PV = -12600$  provided  $PMT = 250$ .

12.56(%) **A1**

**[5 marks]**

*continued...*

Question 3 continued

(c) **EITHER**

Bryan should choose Option A

**A1**

no deposit is required

**R1**

**Note:** Award **R1** for stating that no deposit is required. Award **A1** for the correct choice from that fact. Do not award **R0A1**.

**OR**

Bryan should choose Option B

**A1**

cost of Option A (6923.69) > cost of Option B ( $72 \times 250 - 12\,600 = 5400$ )

**R1**

**Note:** Award **R1** for a correct comparison of costs. Award **A1** for the correct choice from that comparison. Do not award **R0A1**.

**[2 marks]**

(d) real interest rate is  $0.4 - 0.1 = 0.3\%$

**(M1)**

value of other payments  $250 + 250 \times 1.003 + \dots + 250 \times 1.003^{71}$

use of sum of geometric sequence formula or financial app on a GDC

**(M1)**

= 20058.43

value of deposit at the end of 6 years

$1400 \times (1.003)^{72} = 1736.98$

**(A1)**

Total value is (€)21 795.41

**A1**

**Note:** Both **M** marks can awarded for a correct use of the GDC's financial app:

$N = 72$  ( $6 \times 12$ )

$I\% = 3.6$  ( $0.3 \times 12$ )

$PV = 0$

$PMT = -250$

$FV =$

$P/Y = 12$

$C/Y = 12$

**OR**

$N = 72$  ( $6 \times 12$ )

$I\% = 0.3$

$PV = 0$

$PMT = -250$

$FV =$

$P/Y = 1$

$C/Y = 1$

**[4 marks]**

**Total [18 marks]**



4. (a)  $r = \begin{pmatrix} 30 \\ 10 \\ 5 \end{pmatrix} + t \begin{pmatrix} -150 \\ -50 \\ -20 \end{pmatrix}$  **A1A1**

**[2 marks]**

(b) (i) when  $x = 0$ ,  $t = \frac{30}{150} = 0.2$  **M1**

**EITHER**

when  $y = 0$ ,  $t = \frac{10}{150} = 0.2$  **A1**

since the two values of  $t$  are equal the aircraft passes directly over the airport

**OR**

$t = 0.2$ ,  $y = 0$  **A1**

(ii) height =  $5 - 0.2 \times 20 = 1$  km **A1**

(iii) time 13:12 **A1**  
**[4 marks]**

(c) (i)  $5 - 20t = 4 \Rightarrow t = \frac{1}{20}$  (3 minutes) **(M1)**

time 13:03 **A1**

(ii) displacement is  $\begin{pmatrix} 22.5 \\ 7.5 \\ 4 \end{pmatrix}$  **A1**

distance is  $\sqrt{22.5^2 + 7.5^2 + 4^2}$  **(M1)**

= 24.1 km **A1**  
**[5 marks]**

*continued...*

Question 4 continued

(d) **METHOD 1**

time until landing is  $12 - 3 = 9$  minutes

**M1**

height to descend = 4 km

$$a = \frac{-4}{\frac{9}{60}}$$

**M1**

$$= -26.7$$

**A1**

**METHOD 2**

$$\begin{pmatrix} -150 \\ -50 \\ a \end{pmatrix} = s \begin{pmatrix} 22.5 \\ 7.5 \\ 4 \end{pmatrix}$$

**M1**

$$-150 = 22.5s \Rightarrow s = -\frac{20}{3}$$

**M1**

$$a = -\frac{20}{3} \times 4$$


$$= -26.7$$

**A1**

**[3 marks]**

**Total [14 marks]**

5. (a)



**A2**  
[2 marks]

(b) attempt to form an adjacency matrix

$$\begin{pmatrix} 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \end{pmatrix}$$

**M1**

**A1**

[2 marks]

(c) raising the matrix to the power six  
50

**(M1)**

**A1**

[2 marks]

(d) not possible  
because you must pass through B twice

**A1**

**R1**

**Note:** Do not award **A1R0**.

[2 marks]

(e)  $a = 230$ ,  $b = 340$

**A1A1**

[2 marks]

(f)  $A \rightarrow B \rightarrow D \rightarrow E \rightarrow F \rightarrow C \rightarrow A$   
 $90 + 70 + 100 + 210 + 330 + 150$   
(US\$) 950

**(M1)**

**(A1)**

**A1**

[3 marks]

continued...

Question 5 continued

- (g) finding weight of minimum spanning tree **M1**  
 $70 + 80 + 100 + 180 = (\text{US\$}) 430$  **A1**  
 adding in two edges of minimum weight **M1**  
 $430 + 90 + 150 = (\text{US\$}) 670$  **A1**  
**[4 marks]**  
**Total [17 marks]**

6. (a)  $\begin{pmatrix} 0.8 & 0.1 \\ 0.2 & 0.9 \end{pmatrix}$  **M1A1**  
**[2 marks]**  
 (b)  $\begin{vmatrix} 0.8 - \lambda & 0.1 \\ 0.2 & 0.9 - \lambda \end{vmatrix} = 0$  **M1**  
 $\lambda = 1 \text{ and } 0.7$  **A1**  
 eigenvectors  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  and  $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$  **(M1)A1**

**Note:** Accept any scalar multiple of the eigenvectors.

**[4 marks]**

- (c) **EITHER**  
 $P = \begin{pmatrix} 1 & 1 \\ 2 & -1 \end{pmatrix} \quad D = \begin{pmatrix} 1 & 0 \\ 0 & 0.7 \end{pmatrix}$  **A1A1**  
**OR**  
 $P = \begin{pmatrix} 1 & 1 \\ -1 & 2 \end{pmatrix} \quad D = \begin{pmatrix} 0.7 & 0 \\ 0 & 1 \end{pmatrix}$  **A1A1**  
**[2 marks]**

- (d)  $P^{-1} = \frac{1}{3} \begin{pmatrix} 1 & 1 \\ 2 & -1 \end{pmatrix}$  **A1**  
 $\frac{1}{3} \begin{pmatrix} 1 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0.7^n \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1200 \\ 1200 \end{pmatrix}$  **M1A1**  
 attempt to multiply matrices **M1**  
 so in company A, after  $n$  years,  $400(2 + 0.7^n)$  **A1**  
**[5 marks]**

- (e)  $400 \times 2 = 800$  **A1**  
**[1 mark]**

**Total [14 marks]**

7. (a)  $\frac{dv}{dt} = 9.81 - 0.9v$  **M1**
- $\int \frac{1}{9.81 - 0.9v} dv = \int 1 dt$  **M1**
- $-\frac{1}{0.9} \ln(9.81 - 0.9v) = t + c$  **A1**
- $9.81 - 0.9v = Ae^{-0.9t}$  **A1**
- $v = \frac{9.81 - Ae^{-0.9t}}{0.9}$  **A1**
- when  $t = 0$ ,  $v = 0$  hence  $A = 9.81$  **A1**
- $v = \frac{9.81(1 - e^{-0.9t})}{0.9}$
- $v = 10.9(1 - e^{-0.9t})$  **A1**
- [7 marks]**
- (b) **either** let  $t$  tend to infinity, or  $\frac{dv}{dt} = 0$  **(M1)**
- $v = 10.9$  **A1**
- [2 marks]**
- (c)  $\frac{dx}{dt} = y$  **M1**
- $\frac{dy}{dt} = 9.81 - 0.9y^2$  **A1**
- [2 marks]**
- (d)  $x_{n+1} = x_n + 0.2y_n$ ,  $y_{n+1} = y_n + 0.2(9.81 - 0.9(y_n)^2)$  **(M1)(A1)**
- $x = 1.04$ ,  $\frac{dx}{dt} = 3.31$  **(M1)A1**
- [4 marks]**
- (e) 3.3015 **A1**
- [1 mark]**
- (f)  $0 = 9.81 - 0.9(v)^2$  **M1**
- $\Rightarrow v = \sqrt{\frac{9.81}{0.9}} = 3.301511... (= 3.30)$  **A1**
- [2 marks]**

continued...

*Question 7 continued*

- (g) the model found the terminal velocity very accurately, so good approximation **R1**  
intermediate values had object exceeding terminal velocity so not  
good approximation **R1**

**[2 marks]**

**Total [20 marks]**

---