

Applications & Interpretation - 1 Page Formula Sheet

IB Mathematics SL & HL – First examinations 2021

Prior Learning SL & HL	
Area: Parallelogram	$A = bh$, b = base, h = height
Area: Triangle	$A = \frac{1}{2}(bh)$, b = base, h = height
Area: Trapezoid	$A = \frac{1}{2}(a+b)h$, a, b = parallel sides, h = height
Area: Circle	$A = \pi r^2$, r = radius
Circumference: Circle	$C = 2\pi r$, r = radius
Volume: Cuboid	$V = lwh$, l = length, w = width, h = height
Volume: Cylinder	$V = \pi r^2 h$, r = radius, h = height
Volume: Prism	$V = Ah$, A = cross-section area, h = height
Area: Cylinder curve	$A = 2\pi rh$, r = radius, h = height
Distance between two points (x_1, y_1) , (x_2, y_2)	$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$
Coordinates of midpoint	$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$, for endpoints (x_1, y_1) , (x_2, y_2)
Prior Learning HL only	
Solutions of a quadratic equation in the form $ax^2 + bx + c$	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$, $a \neq 0$

Topic 1: Number and algebra - SL & HL	
The n th term of an arithmetic sequence	$u_n = u_1 + (n-1)d$
Sum of n terms of an arithmetic sequence	$S_n = \frac{n}{2}(2u_1 + (n-1)d) = \frac{n}{2}(u_1 + u_n)$
The n th term of a geometric sequence	$u_n = u_1 r^{n-1}$
Sum of n terms of a finite geometric seq.	$S_n = \frac{u_1(r^n - 1)}{r - 1} = \frac{u_1(1 - r^n)}{1 - r}$, $r \neq 1$
Compound interest	$FV = PV \times \left(1 + \frac{r}{100k}\right)^{kn}$ FV is future value, PV is present value, n is the number of years, k is the number of compounding periods per year, $r\%$ is the nominal annual rate of interest
Exponents & logarithms	$a^x = b \Leftrightarrow x = \log_a b$, $a, b > 0$, $a \neq 1$
Percentage error	$\varepsilon = \left \frac{v_A - v_E}{v_E} \right \times 100\%$ v_A = approximate value, v_E = exact value

Topic 1: Number and algebra - HL only	
Laws of logarithms	$\log_a xy = \log_a x + \log_a y$ $\log_a \frac{x}{y} = \log_a x - \log_a y$ $\log_a x^m = m \log_a x$ for $a, x, y > 0$
The sum of an infinite geometric sequence	$S_\infty = \frac{u_1}{1 - r}$, $ r < 1$
Complex numbers	$z = a + bi$
Discriminant	$\Delta = b^2 - 4ac$
Modulus-argument (polar) & Exponential (Euler) form	$z = r(\cos \theta + i \sin \theta) = re^{i\theta} = r \operatorname{cis} \theta$
Determinant of a 2x2 matrix	$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \rightarrow \det A = A = ad - bc$
Inverse of a 2x2 matrix	$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \rightarrow A^{-1} = \frac{1}{\det A} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$
Power formula for a matrix	$M^n = P D^n P^{-1}$, where P is the matrix of eigenvectors and D is the diagonal matrix of eigenvalues

Topic 2: Functions – SL & HL	
Equations of a straight line	$y = mx + c$; $ax + by + d = 0$; $y - y_1 = m(x - x_1)$
Gradient formula	$m = \frac{y_2 - y_1}{x_2 - x_1}$
Axis of symmetry of a quadratic function	$f(x) = ax^2 + bx + c \rightarrow x = -\frac{b}{2a}$
Topic 2: Functions – HL only	
Logistic function	$f(x) = \frac{L}{1 + Ce^{-kx}}$, $L, k, C > 0$

Topic 3: Geometry and trigonometry – SL & HL	
Distance between 2 points (x_1, y_1, z_1) , (x_2, y_2, z_2)	distance $(d) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2}$
Coordinates of the midpoint with endpoints (x_1, y_1, z_1) , (x_2, y_2, z_2)	$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2}\right)$
Volume: Right-pyramid	$V = \frac{1}{3}Ah$, A = base area, h = height
Volume: Right cone	$V = \frac{1}{3}\pi r^2 h$, r = radius, h = height
Area: Cone curve	$A = \pi r l$, r = radius, l = slant height
Volume: Sphere	$V = \frac{4}{3}\pi r^3$, r = radius
Surface area: Sphere	$A = 4\pi r^2$, r = radius
Sine rule	$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
Cosine rule	$c^2 = a^2 + b^2 - 2ab \cos C$ $\cos C = \frac{a^2 + b^2 - c^2}{2ab}$
Area: Triangle	$A = \frac{1}{2}ab \sin C$
Length of an arc	$l = \frac{\theta}{360} \times 2\pi r$ θ = angle in degrees, r = radius
Area of a sector	$A = \frac{\theta}{360} \times \pi r^2$ θ = angle in degrees, r = radius

Topic 3: Geometry and trigonometry – HL only	
Length of an arc	$l = r\theta$ r = radius, θ = angle in radians
Area of a sector	$A = \frac{1}{2}r^2\theta$ r = radius, θ = angle in radians
Identities	$\cos^2 \theta + \sin^2 \theta = 1$ $\tan \theta = \frac{\sin \theta}{\cos \theta}$ $\begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{pmatrix}$ reflection in the line $y = (\tan \theta)x$ $\begin{pmatrix} k & 0 \\ 0 & 1 \end{pmatrix}$ horizontal stretch by scale factor of k $\begin{pmatrix} 1 & 0 \\ 0 & k \end{pmatrix}$ vertical stretch with scale factor of k $\begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}$ centre $(0,0)$ enlargement with scale factor of k $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$, anticlockwise rotation of angle θ about the origin ($\theta > 0$) $\begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$, clockwise rotation of angle θ about the origin ($\theta > 0$)
Magnitude of a vector	$ v = \sqrt{v_1^2 + v_2^2 + v_3^2}$
Vector equ. of a line	$r = a + \lambda b$
Parametric form of the equation of a line	$x = x_0 + \lambda l$, $y = y_0 + \lambda m$, $z = z_0 + \lambda n$
Scalar product	$v \cdot w = v_1 w_1 + v_2 w_2 + v_3 w_3$ $v \cdot w = v w \cos \theta$ where θ is the angle between v and w
Angle between two vectors	$\cos \theta = \frac{v_1 w_1 + v_2 w_2 + v_3 w_3}{ v w }$
Vector product	$v \times w = \begin{pmatrix} v_2 w_3 - v_3 w_2 \\ v_3 w_1 - v_1 w_3 \\ v_1 w_2 - v_2 w_1 \end{pmatrix}$ $ v \times w = v w \sin \theta$ where θ is the angle between v and w
Area of a parallelogram	$A = v \times w $, where v and w form two adjacent sides of a parallelogram

Topic 4: Statistics and probability - SL & HL	
Interquartile range	$IQR = Q_3 - Q_1$
Mean, $\bar{x}$, of a set of data	$\bar{x} = \frac{\sum_{i=1}^n f_i x_i}{n}$, where $n = \sum_{i=1}^k f_i$
Probability of an event A	$P(A) = \frac{n(A)}{n(u)}$
Complementary events	$P(A) + P(A') = 1$
Combined events	$P(A \cup B) = P(A) + P(B) - P(A \cap B)$
Mutually exclusive events	$P(A \cup B) = P(A) + P(B)$
Conditional probability	$P(A B) = \frac{P(A \cap B)}{P(B)}$
Independent events	$P(A \cap B) = P(A)P(B)$
Expected value of a discrete random variable X	$E(X) = \sum x P(X = x)$
Binomial distribution	$X \sim B(n, p)$
Mean ; Variance	$E(X) = np$; $\operatorname{Var}(X) = np(1 - p)$

Topic 4: Statistics and probability – HL only	
Linear transformation of a single random variable	$E(aX + b) = aE(X) + b$ $\operatorname{Var}(aX + b) = a^2 \operatorname{Var}(X)$
Linear combinations of n independent random variables, $X_1, X_2, \dots, X_n$	$E(a_1 X_1 \pm a_2 X_2 \pm \dots \pm a_n X_n) = a_1 E(X_1) \pm a_2 E(X_2) \pm \dots \pm a_n E(X_n)$ $\operatorname{Var}(a_1 X_1 \pm a_2 X_2 \pm \dots \pm a_n X_n) = a_1^2 \operatorname{Var}(X_1) + a_2^2 \operatorname{Var}(X_2) + \dots + a_n^2 \operatorname{Var}(X_n)$
Unbiased estimate of population variance	$s_{n-1}^2 = \frac{n}{n-1} s_n^2$ Sample statistics
Poisson distribution	$X \sim \operatorname{Po}(m)$
Mean ; Variance	$E(X) = m$; $\operatorname{Var}(X) = m$
Transition matrices	$T^n s_0 = s_n$, where s_0 is the initial state

Topic 5: Calculus - SL & HL	
Derivative of x^n	$f(x) = x^n \rightarrow f'(x) = nx^{n-1}$
Integral of x^n	$\int x^n dx = \frac{x^{n+1}}{n+1} + C$, $n \neq -1$
Area enclosed by a curve and the x-axis	$A = \int_a^b y dx$, where $f(x) > 0$
The trapezoidal rule	$\int_a^b y dx \approx \frac{b-a}{n} h \left((y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1}) \right)$ where $h = \frac{b-a}{n}$

Topic 5: Calculus – HL only	
Derivative of $\sin x$	$f(x) = \sin x \rightarrow f'(x) = \cos x$
Derivative of $\cos x$	$f(x) = \cos x \rightarrow f'(x) = -\sin x$
Derivative of $\tan x$	$f(x) = \tan x \rightarrow f'(x) = \frac{1}{\cos^2 x}$
Derivative of e^x	$f(x) = e^x \rightarrow f'(x) = e^x$
Derivative of $\ln x$	$f(x) = \ln x \rightarrow f'(x) = \frac{1}{x}$
Chain rule	$y = g(u)$, $u = f(x) \rightarrow \frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$
Product rule	$y = uv \rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$
Quotient rule	$y = \frac{u}{v} \rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
Standard integrals	$\int \frac{1}{x} dx = \ln x + C$ $\int \sin x dx = -\cos x + C$ $\int \cos x dx = \sin x + C$ $\int \frac{1}{\cos^2 x} dx = \tan x + C$ $\int e^x dx = e^x + C$
Area enclosed by a curve and x or y-axes	$A = \int_a^b y dx$ or $A = \int_a^b x dy$
Volume of revolution about x or y-axes	$V = \int_a^b \pi y^2 dx$ or $V = \int_a^b \pi x^2 dy$
Acceleration	$a = \frac{dv}{dt} = \frac{d^2 s}{dt^2} = v \frac{dv}{ds}$
Distance; Displacement travelled from t_1 to t_2	$\text{dist} = \int_{t_1}^{t_2} v(t) dt$; $\text{disp} = \int_{t_1}^{t_2} v(t) dt$
Euler's method	$y_{n+1} = y_n + h \times f(x_n, y_n)$; $x_{n+1} = x_n + h$ where h is a constant (step length)
Euler's method for coupled systems	$x_{n+1} = x_n + h \times f_1(x_n, y_n, t_n)$ $y_{n+1} = y_n + h \times f_2(x_n, y_n, t_n)$ $t_{n+1} = t_n + h$ where h is a constant (step length)
Exact solution for coupled linear differential equations	$x = Ae^{\lambda_1 t} p_1 + Be^{\lambda_2 t} p_2$



www.revisionvillage.com

IB Maths Exam Questionbanks

IB Maths Learning Videos

IB Maths Past Paper Video Solutions

IB Maths Practice Exams (Full Length)

IB Maths Practice Exams by Topic

IB Maths Practice Exams by Sub-Topic

Voted #1 IB Maths Resource Online & Used by 250,000+ IB Students & Teachers Worldwide